

ACITAIM 25mg Capsules

ACITRETIN

ایسی ٹیم ۲۵ ملی گرام کپسولز
ایسی ٹیمین

COMPOSITION:

Each capsule contains:

Acitretin 25mg

(USP Specifications)

INDICATIONS:

Acitaim (Acitretin) is a second-generation retinoid indicated for the treatment of severe psoriasis in adults g. Plaque Psoriasis, Guttate Psoriasis, Inverse Psoriasis, Pustular Psoriasis & Erythrodermic Psoriasis.

OTHER

Acitaim is sometimes also used to treat Darier's disease, palmoplantar pustulosis, lichen sclerosus elatrophicus of the vulva, plamoplantar lichen nitidus, and lichen planus, it is also used to trout lamellar ichthyosis, sjogren Larsson syndrome, and bullous and nonbullous Ichthyosiform erythroderma.

Dosage:

Acitaim therapy should be initiated at 25 to 50mg per day, given as a single dose with the main meal. Maintenance doses of 25 to 50mg per day may be given dependent upon an individual patient's response to initial treatment. Relapses may be treated as outlined for initial per day may be given dependent upon an Individual patient's response to initial treatment.

Contraindications and warnings:

For Female Patients:

Acitaim must not be used by females who are pregnant or who may become pregnant during therapy or at any time for at least 3 years after discontinuation of treatment. Acitaim also must not be used by females of reproductive potential who may be used 2 effective forms of contraception (birth, control) simultaneously, for at least one month before during and for at least three years after treatment. Two effective forms of contraception (birth control) are to be used simultaneously even when one is a hormonal contraceptive. Patients should not self-medicate with St. John's Wort because of possible interaction with hormonal contraceptives. Prescribers must obtain negative results for 2 Pregnancy tests before initiating treatment with Acitaim. The first test is a screening test, the second is a confirmation test done during the first 5 years of the menstrual period immediately preceding Acitaim therapy. For Patients with menorrhoea, the second lest should be done at least 11 days after the last act of unprotected sexual intercourse. Timing of Pregnancy Testing throughout the treatment course should be monthly or individualized based on the prescriber's clinical judgement. Females must sign a Patient Information/Consent about the risks of birth defects. Acitretin is a metabolite of etretinate and major tetsl abnormalities have been reported with both drugs. Acitretin can interact with ethanol to form etretinate Therefore, females of reproductive potential must not ingest ethanol during treatment and for 2 months after cessation of treatment. Before prescribing. Please see Complete late pregnancy warning in the accompanying complete product information. Females who have undergone treatment with tretinate must continue to follow the contraception

requirements for etretinate

For male and female Patients:

Do not donate blood while taking Acitaim and for 3 years after treatment. Acitaim may cause liver damage. Tell your doctor if you have or have ever had liver disease.

Common side effects:

Common side effects include: cheilitis, rhinitis, dry mouth, arthralgia, spiral hyperostosis, alopecia, skin peeling dry skin nail disorder pruritus, erythematous rash, Less frequent, but potentially serious, adverse events include hepatotoxicity pancreatitis as well as hyperostosis, bed headaches, nausea, vomiting Plumed vision, decreased vision in the dark depression yellowing of your our skin great thirst or hunger, shortness of breath, aftersation in lipids with possible cardiovascular affects

Presentation

Acitaim Capsules are available in Alu-Alu blister pack of 3 x 10

PRECAUTION:

- Use as prescribed by the physician.
- Keep out of the reach of children.
- Store at room temperature.
- Protect from direct sunlight, heat & moisture.



Manufactured by:

Aims Pharmaceuticals

Plot # 291, Industrial Triangle
Kahuta Road, Islamabad - Pakistan.
www.aimspharmaceuticals.com

ہدایات:

دوا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بچوں کی پہنچ سے دور رکھیں۔

گرمی اور حرارت سے محفوظ رکھیں۔

سورج کی روشنی، گرمی اور نمی سے محفوظ رکھیں۔