

Clobecin Ointmet

(Clobetasole Propionate USP)

کلو بیسین اسٹیمٹ

Comosition:

Clobetasol Propionate 0.05w/w (USP Specs)

DESCRIPTION

Clobetasole Propionate is a synthetic corticosteroid for topical dermatological uses

INDICATIONS:

It is topical corticosteroid used in short courses for psoriasis excluding wide spread other plaque psoriasis recalcitrant aczema Lichen planus Discold lupus erythemantu and other skin condition which donot respond satisfactorily to less active steroids.

DOSAGE & ADMINISTRATION:

Apply sparingly to the affected area once or twice daily until improvement occurs. Discontinue therapy when control is achieved. Treatment should not continue for more than four weeks without reviewing the patient's condition. Repeated shorts courses may be used to control exacerbations. Not recommended for long-term use Occlusive Apply sparingly to the affected area once or twice daily until improvement occurs. Dressing is recommended in more resistant cases.

CONTRA INDICATION:

Rosacea acne vulgaris, perioral dermatitis, primary cutaneous Viral infections infected skin lesions caused by infection with fungi (e. candidiasis, tinea) or bacteria y e.g. impetigo of dermatosis in children under 1 year of age, including dermatitis and napkin eruptions.

PRECAUTIONS & WARNINGS:

Avoid prolong application to the face. Ensure preparation does not enter the eyes as glaucoma may result, infection may be masked. Clobecin should not be used for long-term especially in children, due to the risk of adrenal suppression. Atrophic changes of the skin may occur after prolonged treatment. It should be applied to the eyelids with care psoriatic patients should be treated under supervision. Infected lesions may require systemic antibiotics therapy. if infection spread. Clobecin should be discontinued. Occlusive dressings may encourage bacterial infection.

Topical steroid should not be used extensively in pregnancy

ADVERSE EFFECTS:

Prolonged use of large amounts or treatment extensive areas can cause features of hypercriticism. Prolonged use may also cause atrophy of skin e.g. thinning, striae and dilation of superficial blood vessels. Pigmentation changes hypertrichosis, hypersensitivity, exacerbations of symptoms may occur, treatment of psoriasis (or its withdrawal) may provoke the pustular form of the disease.

PRESENTATIONS: Packing

PRECAUTION:

- Use as prescribed by the physician.
- Keep out of the reach of children.
- Store at room temperature.
- Protect from direct sunlight, heat & moisture.

ہدایات:

دوا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بچوں کی پہنچ سے دور رکھیں۔

کمرے کے درجہ حرارت پر رکھیں۔

سورج کی روشنی، گرمی اور نمی سے محفوظ رکھیں۔



Manufactured by:

Aims Pharmaceuticals

Plot # 291, Industrial Triangle
Kahuta Road, Islamabad - Pakistan.
www.aimspharmaceuticals.com