

ROTIN

10 mg
TABLETS
(Loratadine)

روٹین
لوراناڈین
گولیاں

COMPOSITIONS:

Each tablet contains: Loratidine.....10mg

DESCRIPTION: Loratidine is a potent and long acting antihistamine with selective peripheral H₁-receptor antagonist activity.

PHARMACOKINETICS: Loratidine is well absorbed and extensively metabolized. It reaches in measurable levels in plasma within 15 minutes and peak levels are achieved with in 1 hour after single oral dose while its elimination half-life is 18-24 hours. It can be given with or without food, since food doses not affect the pharmacokinetics of Loratidine. **Renal insufficiency & Hepatic Impairment:** Results of Loratidine studies in patients with chronic renal insufficiency and patients with alcoholic liver Disease have shown that these impairments have no meaningful effect on the pharmacokinetics of Loratidine.

INDICATIONS: ROTIN (Loratidine) tablets are indicated for the relief of symptoms associated with allergic rhinitis such as sneezing, nasal discharge (rhinorrhea), itching, ocular itching & burning chronic urticaria and other dermatological conditions associated with allergy.

USUAL DOSAGE: Adults and children above 12 years: 1 tablet ROTIN (Loratidine 10mg) **Children 2-12 years:** Less than 30kg: ROTIN (Loratidine 10mg) once daily. More than 30kg: 1 tablets of ROTIN) Loratidine 10mg) daily OR as advised by the physician.

DRUG INTERACTIONS: Loratidine doses not potentiate the effects of alcohol. Anti-histamines should be discontinued about 1 days prior to skin testing procedures, since these drugs may prevent or diminish otherwise positive reactions to dermal reactivity indicators.

POSSIBLE ADVERSE EFFECTS: Loratidine has no clinically significant sedative or anti-cholinergic properties since fatigue, sedation, headache, nausea and dry mouth were rarely reported.

CONTRAINDICATIONS: Loratidine is contraindicated in patients who have shown hypersensitivity or idiosyncrasy to any of its components.

WARNINGS: Loratidine should be prescribed with caution in pregnancy and lactation since its safety is not established in such conditions.

Pack size

Rotin is available as:

Loratidine 10 mg=1x10's Tab/pack.

PRECAUTION:

- Use as prescribed by the physician.
- Keep out of the reach of children.
- Store at room temperature.
- Protect from direct sunlight, heat & moisture.

ہدایات:

دواؤں کی زیادتی کے مطابق استعمال کریں۔

بچوں کی پہنچ سے دور رکھیں۔

کمرے کے درجہ حرارت پر رکھیں۔

سورج کی روشنی، گرمی اور نمی سے محفوظ رکھیں۔



Manufactured by:

Aims Pharmaceuticals

Plot # 291, Industrial Triangle
Kahuta Road, Islamabad - Pakistan.
www.aimspharmaceuticals.com