

Vosul 
(Levosulpiride) Tablets
25mg
50mg

ووسل 
(لیووسلپیراید)

Composition:

VOSUL-25mg Tablets

Each tablet contains Levosulpiride25mg

VOSUL-50 mg Tablets:

Each tablet contains Levosulpiride50mg

Classification:

Levosulpiride is a substituted benzamide, chemically is S-(-)-N(1-ethylpyridine-2II-methyl 1-5-) sulfamoyl benzamide.

Therapeutic Classifications:

Atypical neuroleptic with srereo selective antidopaminergic action.

Properties:

The biochemical, pharmacological and clinical data obtained from the two sulphiride isomers indicated that the antidopaminergic action, both on a central and peripheral level, is due to the levogyre enantiomer (Levosulpiride) VOSUL exhibits a clear antidepressive, anticemesihopatic uninhibiting a lesocializing action. Due to its hallucinolytic and delriolytic action. it is effective against acute and chronic schizophrenia For depression. VOSUL is administered at lower dosages than those used on patients with schizophrenia. It is suggested that these low dosages involve a preferential action on the plesynaptic DA-Receptors, thus producing a functional increase of dopaminic transmission and stimulating motoric and psychic action, The antipsychotic effects of VOSUL generated at high dosages may be due to a blocking of the post synaptic DA-receptors although these effects are not as characteristic as those produced by classic neuroleptic agents .VOSUL, exerts its action at dosages which are 2-3 times lower than those of the racemic molecule, which entails a better tolerability and a reduced occurrence of collateral effects, Sedation antimuscannic effects, hypotension and extrapyramidal disorders are less of a problem than with most other neuroleptics

Pharmacokinetics:

After oral administration the bioavailability of L-sulpiride is about 30% Peak-plasma concentration occurs after about 3 hours and it has plasma half life of about 9-7 hours. It is mostly eliminated by the kidneys in the urine.

Indications:

VOSUL is indicated in the treatment of following disorders.

- Endogenous and reactive depressive states.
- sometic disturbances
- Acute and chronic schizophrenia especially recommended for negative syptoms (after disorders).

Dosage:

- Endogenous and reactive depressive states/somatic disturbances, For Adults 2-3 tablets of 50mg daily
- Acute and chronic schizophrenia: For Adults 4-6 tablets of 50mg daily.

Maintenance therapy. The dosages may be progressively lowered when treating aged patients the dosage must be carefully established by the physician who must consider a possible reduction in the dosage mentioned above.

Contraindications:

As with other neuroleptics, levosulpiride is also contraindicated in patients with hyperprolactinemic tendency mammary dysplasias and malignant mastopathies

Precautions:

VOSUL is to be applied with caution on epilepsies, manic states, manic stages of manic-depressive psychoses cases of insufficiency

Warnings:

As with other neuroleptics the association of levosulpiride with typical neuroleptics demands particular medical care and attention, In case of potentially fatal complex of symptoms, known as malignant Neuroleptic Syndrome (hyperpyrexia, muscular rigidity akinesia, vegetative disturbances: irregularities of pulse and arterial pressure regulation, anhythmias. alterations of the conscious conditions or numbness and coma) Under such circumstances the treatmet may be suspended should be continued if considered essential and patients must be carefully monitored.

Drug interaction:

As with other neuroleptics or substituted benzamide the concomitant use of levosulpiride with therapeutic doses of sucralfate or an antacid containing aluminium and magnesium hydride may reduce mean oral bioavailability of levosulpiride.

Side Effects:

Numbness a sense of vertigo and asynchronisms may occur patients under treatment must be made aware of these risks in order to avoid driving vehicles and attending to operations demanding total attention due to their potential dangers. In very rare cases psychomotoric excitement, neurovegetative disturbances and allergic phenomena have been observed, Isolated cases of an Increase in body weight have reported following prolonged treatment. All these effects are of a modest and reversible levosulpiride on the functionality of the libida observed on particular occasions are reconducibile to a reversible effects of levosulpiride on the functionality of the hypothalamus pituitary gonadic axis which are similar to that known in many neuroleptics.

Precautions

- Keep out of the reach of children .
- Store at room temperature.
- Protect from direct sunlight, heat & moisture.

Pack Size

VOSUL 25mg is available in a box of 20 tablets
VOSUL 50mg is available in a box of 20 tablets

ہدایات:

دوا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ٹھنڈی اور خشک جگہ پر رکھیں۔

بچوں کی پہنچ سے دور رکھیں۔



Manufactured by:

Aims Pharmaceuticals

Plot # 291, Industrial Triangle
Kahuta Road, Islamabad - Pakistan.
www.aimspharmaceuticals.com